



Europäisches Patentamt
European Patent Office
Office européen des brevets

Publication number:

**0 330 026
A1**

EUROPEAN PATENT APPLICATION

Application number: 89102376.4

Int. Cl. 4: **C07D 211/26 , C07D 401/12 ,
C07D 405/12 , C07D 409/12 ,
A61K 31/445**

Date of filing: 11.02.89

Priority: 15.02.88 JP 32339/88
11.05.88 JP 114169/88

Date of publication of application:
30.08.89 Bulletin 89/35

Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE


Applicant: **TAKEDA CHEMICAL INDUSTRIES,
LTD.**
3-6, Doshomachi 2-chome Chuo-ku
Osaka(JP)

Inventor: **Goto, Glichi**
6-11, Kofudai 5-chome Toyono-cho
Toyono-gun Osaka 563-01(JP)
Inventor: **Nagaoka, Akinobu**
45-8, Daiwahigashi 1-chome
Kawanishi Hyogo 666-01(JP)

Representative: **von Kreisler, Alek,**
Dipl.-Chem. et al
Patentanwälte Von Kreisler-Selting-Werner
Deichmannhaus am Hauptbahnhof
D-5000 Köln 1(DE)

Piperidinoalkyl derivatives of carboxylic acid amides.

The present invention relates to unsaturated carboxylic acid amide derivatives of the formula



wherein ring A stands for an optionally substituted aromatic ring; R¹ stands for a hydrogen atom or an optionally substituted hydrocarbon residue or forms an optionally substituted carbocyclic ring with the adjacent group -CH=C- together with two carbon atoms constituting the ring A; R² stands for a hydrogen atom, an optionally substituted hydrocarbon residue or an optionally substituted acyl group; R³ stands for an optionally substituted hydrocarbon residue; and n denotes an integer ranging from 2 to 6, and salts thereof, as well as the production thereof.

The compounds of the present invention act on the central nervous system of mammals and has a strong anti-cholinesterase activity, which can be used for the prophylaxis and therapy of, for example, senile dementia, Alzheimer's diseases, Huntington's chorea, et., and are useful as medicines.

EP 0 330 026 A1

Piperidinoalkyl Derivatives of Carboxylic Acid Amides


This invention relates to novel unsaturated carboxylic acid amide derivatives useful as Pharmaceuticals, especially as agents for improving cerebral functions in senile dementia, Alzheimer's disease, etc.

With the increase of aged population, a variety of compounds having actions of improving cerebral functions have been suggested. Among them, physostigmine, an anticholinesterase agent, has been found to have an action of improving cerebral functions.

Physostigmine, however, has such drawbacks as a short duration of action and high toxicity.

The object of the present invention is to provide compounds with longer action and with lower toxicity, as compared with known compounds improving cerebral functions.

The present inventors succeeded in creating unsaturated carboxylic acid amide derivatives represented by the general formula (I):



wherein ring A stands for an optionally substituted aromatic ring; R¹ stands for a hydrogen atom or an optionally substituted hydrocarbon group or forms an optionally substituted carbocyclic ring with the adjacent group -CH=C- together with two carbon atoms constituting the ring A; R² stands for a hydrogen atom, an optionally substituted hydrocarbon group or an optionally substituted acyl group; R³ stands for an optionally substituted hydrocarbon group; and n denotes an integer ranging from 2 to 6, or salts thereof, and found that these compounds have a strong cholinesterase antagonizing action as well as a potent action of improving cerebral functions.

The present invention relates to compounds represented by the formula (I) or salts thereof, methods of preparing them and anticholinesterase agents containing them as well as agents containing them for improving cerebral functions.

In the above formula (I), examples of "hydrocarbon group" of "optionally substituted hydrocarbon group" represented by R¹, R² and R³ include those in the form chain-like, cyclic, saturated or unsaturated group and combinations thereof. Examples of chain-like saturated hydrocarbon groups in chain form include straight-chain or branched alkyl groups of 1 to 11 carbon atoms (e.g. methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, tert-butyl, n-pentyl, n-hexyl).

Examples of chain-like unsaturated hydrocarbon groups in chain form include straight-chain or branched C₂-4 alkenyl e.g. vinyl, allyl, 2-butenyl, and C₂-4 alkynyl groups (e.g. propargyl, 2-butylnyl).

Examples of cyclic saturated hydrocarbon groups include C₃-7 monocyclic cycloalkyl (e.g. cyclobutyl, cyclopentyl, cyclohexyl), and C₈-14 cross-linked cyclic saturated hydrocarbons, e.g. bicyclooctyls such as bicyclo[3,2,1]oct-2-yl, bicyclononyls such as bicyclo[3,3,1]nonan-2-yl, and tricyclodecyls such as adamantan-1-yl. Cyclic unsaturated hydrocarbon groups include phenyl group, naphthyl group, etc.

Examples of substituents of these hydrocarbon groups include such groups as halogen atoms (e.g. chlorine, bromine, iodine), nitro, nitrile, hydroxy, C₁-4 alkoxy (e.g. methoxy, ethoxy, propoxy, butyloxy, isopropoxy), C₁-4 alkylthio (e.g. methylthio, ethylthio, propylthio, isopropylthio, butylthio), amino, mono or di-C₁-4 alkyl substituted amino (e.g. methylamino, ethylamino, propylamino, dimethylamino, diethylamino), C₁-4 alkoxy carbonyl (e.g. methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isobutoxycarbonyl), hydroxycarbonyl, C₁-5 alkyl carbonyl (e.g. methylcarbonyl, ethylcarbonyl, butylcarbonyl, cyclohexylcarbonyl), carbamoyl, mono- or di-C₁-4 alkyl substituted carbamoyl (e.g. methylcarbamoyl, ethylcarbamoyl, propylcarbamoyl, butylcarbamoyl, diethylcarbamoyl, dibutylcarbamoyl), phenyl, naphthyl, phenoxy, benzoyl, phenoxycarbonyl, phenyl C₁-4 alkyl carbamoyl and phenyl carbamoyl which may have 1 to 4 substituents and adamantan-1-yl. Substituents on the phenyl groups or naphthyl groups are exemplified by C₁-4 alkyl groups such as methyl, ethyl, propyl, butyl, isopropyl, etc. The phenyl groups can optionally have 1 to 4 substituents (substituents on the phenyl group are exemplified by C₁-4 alkyl group such as methyl, ethyl, propyl, butyl, isopropyl, etc., halogen such as chlorine, bromine, iodine, etc., hydroxy, benzyloxy, amino, mono- or di-C₁-4 alkyl-substituted amino, nitro, C₁-4 alkoxy carbonyl, etc.).

The preferred number of substituents of these hydrocarbon groups is 1 to 3.

The aromatic ring represented by the ring A includes aromatic cyclic compounds such as aromatic monocyclic hydrocarbon, aromatic condensed polycyclic hydrocarbon or aromatic heterocyclic ring. The aromatic monocyclic hydrocarbon is exemplified by benzene, and the aromatic condensed polycyclic hydrocarbon is exemplified by naphthalene, anthracene, etc. The aromatic heterocyclic ring includes 5- to 6-membered heterocyclic rings containing 1 to 4 hetero atoms selected from nitrogen atom, oxygen atom and sulfur atom. These aromatic heterocyclic rings are exemplified by thiophene, furan, pyrazole, thiazole, isothiazole, oxazole, isoxazole, imidazole, triazole, tetrazole, pyridine, pyrimidine, pyridazine, etc.

Among these heterocyclic rings, thiophene, furan, pyridine, etc., containing one hetero atom, are especially preferred.

In the above formula (I), R^1 may form a carbocyclic ring together with the adjacent group $-CH=C-$ and two carbon atoms constituting the ring A. As such carbocyclic rings, 5-to 7-membered rings are especially desirable. As such carbon rings those having no aromaticity are especially desirable. The condensed ring formed by such a carbon ring as above and the ring A is exemplified by indene, 1,2-dihydronaphthalene, 6,7-dihydro-5H-benzocycloheptene, 5,6,7,8-tetra-hydrocyclooctene, 4,5-dihydrobenzo[b]thiophene, 4,5-dihydroisobenzofuran, 7,8-dihydroquinoline, 7,8-dihydroisoquinoline, etc. As substituents on these carbon rings are used C_{1-4} alkyl groups such as methyl, ethyl, propyl, isopropyl, butyl, etc., halogen such as chlorine, bromine, iodine, hydroxyl group, C_{1-4} alkyloxy groups such as methoxy, ethoxy, propyloxy, isopropyloxy, butyloxy, amino, mono- or di- C_{1-4} alkyl-substituted amino such as methylamino, dimethylamino, etc., nitro, cyano, C_{1-4} alkoxycarbonyl such as methoxycarbonyl, etc. The number of substituents on these carbon rings is preferably 1 to 3.

The acyl group represented by R^2 in the above formula (I) is exemplified by carboxylic acid acyl, carbamic acid acyl, sulfonic acid acyl, substituted hydroxycarboxylic acid acyl etc. The acyl groups may optionally have substituents.

Examples of the carboxylic acid acyl include C_{1-6} alkylcarbonyl e.g. acetyl, propionyl, butyryl, valeryl, hexanoyl, isobutyryl, isovaleryl, etc., C_{3-8} cycloalkylcarbonyl e.g. cyclopentanoyl, cyclohexanoyl, etc., C_{3-8} cycloalkyl- C_{1-6} alkylcarbonyl e.g. cyclopentylacetyl, etc., C_{2-6} alkenyl or alkynylcarbonyl e.g. acryloyl, crotonyl, 2-pentenoyl, 4-pentynoyl, 2-hexenoyl, 3-hexenoyl, 2,4-hexadienoyl, etc., arylcarbonyl e.g. benzoyl, naphthoyl, etc.

As the carbamic acid acyl, use is made of, for example, carbamoyl, mono- or di-substituted carbamoyl. Examples of the mono- or di-substituted carbamoyl include mono- or di- C_{1-4} alkylcarbamoyl such as methylcarbamoyl, ethylcarbamoyl, propylcarbamoyl, butylcarbamoyl, dimethylcarbamoyl, diethylcarbamoyl, dipropylcarbamoyl, etc., mono- or di- C_{3-6} alkenyl carbamoyl such as allylcarbamoyl, 3-butenylcarbamoyl, 4-pentenylcarbamoyl, diallylcarbamoyl, etc., alkynylcarbamoyl or aromatic carbamoyl groups such as phenylcarbamoyl, naphthylcarbamoyl, diphenylcarbamoyl, etc.

Examples of the sulfonic acid acyl groups include inorganic sulfonyl groups such as sodium sulfonyl, C_{1-6} alkylsulfonyl such as methylsulfonyl, ethylsulfonyl, propylsulfonyl, butylsulfonyl, etc., C_{2-6} alkenyl- or alkynyl-sulfonyl such as allylsulfonyl, 2-methyl-2-propenylsulfonyl, etc., aromatic sulfonyl such as phenylsulfonyl, naphthalenesulfonyl, etc.

Examples of the substituted hydroxycarboxylic acid acyl groups include C_{1-6} alkyloxycarbonyl such as methyloxycarbonyl, ethyloxycarbonyl, tert-butyloxycarbonyl, hexyloxycarbonyl, etc., C_{3-8} cycloalkyloxycarbonyl such as cyclopentyloxycarbonyl, cyclohexyloxycarbonyl, etc., cycloalkyl-alkyloxycarbonyl such as cyclopentylmethyloxycarbonyl, etc., C_{2-7} alkenyl- or alkynyl-oxycarbonyl such as allyloxycarbonyl, crotyloxycarbonyl, 2-penten-1-oxycarbonyl, etc., or aromatic or aromatic aliphatic hydroxycarbonyl such as phenyloxycarbonyl, or phenyl- C_{1-2} alkoxycarbonyl such as benzyloxycarbonyl, etc.


As the substituents where these acyl groups have further substituents, these substituents can be groups exemplified by the above-mentioned substituents of hydrocarbon residues. The preferable number of such substituents is 1 to 3.

Referring to the stereochemistry at the site of the double bond shown by $-CH=C-$ of the above formula (I), either the E-isomer or Z-isomer or a mixture thereof may be usable.

As the substituents which the ring A in the above formula (I) may optionally have (one of the substituents on the ring A is sometimes shown as X), there can be used, for example, a C_{1-4} alkyl (e.g. methyl, ethyl, propyl, butyl, etc.), a halogen atom (e.g. chlorine, bromine, iodine), nitro, cyano, hydroxy, a C_{1-4} alkoxy (e.g. methoxy, ethoxy, propyloxy, butyloxy, isopropyloxy, etc.), a C_{1-4} alkylthio (e.g. methylthio, ethylthio, propylthio, isopropylthio, butylthio, etc.), amino, a mono- or di- C_{1-4} alkyl-substituted amino (e.g. methylamino, ethylamino, propylamino, dimethylamino, diethylamino, etc.), C_{1-4} alkylcarbonylamino (e.g. acetylamino, propionylamino, butyrylamino, etc.), C_{1-4} alkylsulfonylamino e.g. methylsulfonylamino, ethylsulfonylamino, propylsulfonylamino, etc.), C_{1-4} alkoxycarbonyl (e.g. methoxycarbonyl, ethoxycarbonyl,

propoxycarbonyl, isobutoxycarbonyl, etc.), hydroxycarbonyl, C₁₋₆ alkylcarbonyl (e.g. methylcarbonyl, ethylcarbonyl, butylcarbonyl, cyclohexylcarbonyl, etc.), carbamoyl, mono- or di-C₁₋₄ alkyl-substituted carbamoyl (e.g. methylcarbamoyl, ethylcarbamoyl, propylcarbamoyl, butylcarbamoyl, diethylcarbamoyl, dibutylcarbamoyl, etc.), C₁₋₆ alkylsulfonyl (e.g. methylsulfonyl, ethylsulfonyl, propylsulfonyl, cyclopentylsulfonyl, cyclohexylsulfonyl, etc.), phenyl, phenoxy, benzoyl, phenoxycarbonyl, phenyl C₁₋₄ alkylcarbamoyl, phenylcarbamoyl, phenyl C₁₋₄ alkylcarbonylamino, benzoylamino, phenyl-C₁₋₄ alkylsulfonyl, phenylsulfonyl, phenyl C₁₋₄ alkylsulfonylamino and phenylsulfonylamino group, which may have 1 to 4 further substituents. The phenyl groups in these substituents can be further substituted by C₁₋₄ alkyl groups such as methyl, ethyl, propyl, butyl, isopropyl, etc., halogen such as chlorine, bromine, iodine, hydroxy, benzyloxy, amino, mono- or di-C₁₋₄ alkyl-substituted amino such as methylamino, dimethylamino, etc., nitro, C₁₋₄ alkoxycarboxy groups. The number of substituents on the ring A is preferably 1 to 3. Ring A is preferably unsubstituted or has one or two of the above substituents.

When R¹ forms an optionally substituted carbocyclic ring with the adjacent group -CH=C- together with two carbon atoms constituting the ring A, compound (I) can be represented by the formula (I'):



wherein A' is a ring defined in the same way as ring A above; X' is hydrogen, C₁₋₄ alkyl, halogen, hydroxy, C₁₋₄ alkoxy, amino, mono- or di-C₁₋₄ alkylamino, cyano or C₁₋₄ alkoxycarbonyl etc.; a is an integer from 1 to 3, and other the symbols are defined above.

Referring to preferable embodiments of the compounds shown by the above formula (I), as the ring A, benzene, pyridine, furan and thiophene can be used, and benzene and pyridine are especially preferable.

Preferable examples of R¹ include a hydrogen atom, C₁₋₆ alkyl group such as methyl, ethyl, propyl, etc., a substituted or unsubstituted phenyl group, or R¹ can also preferably be a group forming 1,2-dihydronaphthalene or 6,7-dihydro-5H-benzocycloheptane together with the adjacent group -CH=C- and two carbon atoms of the ring A. Especially preferred R¹ is a hydrogen atom or phenyl or where 6,7-dihydro-5H-benzocycloheptene is formed in combination with the adjacent double bonded group -CH=C- and the ring A.

Preferable examples of R² include a hydrogen atom, C₁₋₆ alkyl such as methyl, ethyl, propyl, etc., optionally substituted phenyl, C₁₋₆ alkylcarbonyl such as acetyl, propionyl, butyryl, etc. or arylcarbonyl such as benzoyl, especially hydrogen atom, C₁₋₆ alkylcarbonyl or benzoyl.

As R², an optionally C₁₋₃ alkyl substituted aromatic hydrocarbon, e.g. phenyl, naphthyl or phenyl-C₁₋₃ alkyl can be used, benzyl is especially preferred.

Preferable examples of the substituents (X) on the ring A include a C₁₋₄ alkyl group such as methyl, ethyl, propyl, etc., halogen atom such as chlorine, bromine, etc., nitro, cyano, a C₁₋₄ alkoxy group such as methoxy, ethoxy, propoxy, etc., an optionally substituted phenoxy C₁₋₄ alkylcarbonylamino such as acetaminato, propionylamino, C₁₋₄ alkylsulfonylamino such as methylsulfonylamino, ethylsulfonylamino, etc., phenyl C₁₋₄ alkylsulfonylamino such as benzylsulfonylamino, etc., optionally substituted phenylsulfonylamino, C₁₋₄ alkylcarbonyl such as acetyl, propionyl, butyryl, etc., C₁₋₄ alkoxycarbonyl such as methoxycarbonyl, ethoxycarbonyl, butoxycarbonyl, etc., optionally substituted phenoxycarbonyl, optionally substituted benzoyl, carbamoyl, mono- or di-C₁₋₄ alkyl-substituted carbamoyl such as methylcarbamoyl, ethylcarbamoyl, butylcarbamoyl, etc., optionally substituted phenylcarbamoyl, optionally substituted C₁₋₆ alkylthio such as methylthio, ethylthio, propylthio, etc., optionally substituted phenyl-C₁₋₄ alkylthio such as benzylthio, phenethylthio, etc., optionally substituted C₁₋₆ alkylsulfinyl such as methylsulfinyl, ethylsulfinyl, propylsulfinyl, etc., optionally substituted phenyl-C₁₋₆ alkylsulfinyl such as benzylsulfinyl, phenethylsulfinyl, etc., C₁₋₆ alkylsulfonyl such as methylsulfonyl, propylsulfonyl, cyclohexylsulfonyl, etc., optionally substituted phenyl-C₁₋₄ alkylsulfonyl, phenyl-C₁₋₄ alkyl such as optionally substituted phenylsulfonyl, optionally substituted phenyl, phenyl-C₁₋₄ alkyl such as optionally substituted benzyl, etc. Among them, C₁₋₄ alkyl, halogen nitro, cyano, acetaminato, C₁₋₄ alkoxy, optionally substituted phenyl, optionally substituted benzyl, optionally substituted benzoyl, optionally substituted benzoylamino, optionally substituted C₁₋₆ alkylsulfonyl, optionally substituted benzylsulfonyl, optionally substituted phenylsulfonylamino, optionally substituted

benzylsulfonylamino, optionally substituted phenylcarbamoyl, methoxycarbonyl, diethoxycarbonyl, etc. are especially preferable.

The substituent X on ring A which contains a phenyl group therein is referred to herein also as group Q. Thus, group Q includes a phenyl, benzyl, benzoyl, benzoylamino, benzylsulfonyl, phenylsulfonylamino, benzylsulfonylamino, or phenylcarbamoyl etc. The group Q may be optionally substituted with 1 to 3 of C₁₋₄ alkyl, phenyl, halogen, hydroxy, benzyloxy, amino, mono- or di-C₁₋₄ alkylamino, nitro or C₁₋₄ alkoxy carbonyl etc.

As X, electron attractive groups are especially preferable among the above-mentioned groups.

The ring having no substituent is also preferable.

As n are preferable 2,3 and 4.

Especially preferred are compounds (I) wherein the ring A is benzene or pyridine; R¹ is a hydrogen atom, methyl or phenyl group or forms 6,7-dihydro-5H-benzocycloheptene or 1,2-dihydronaphthalene together with the adjacent group -CH=C- and the ring A; R² is an acetyl or propionyl group, R³ is a benzyl group; n is 2, 3 or 4; and the ring A is unsubstituted or substituted with a nitro group or acylamino group.


The compound (I) of the present invention may form an acid addition salt, especially a physiologically acceptable acid addition salt. As these salts, mention is made of, for example, salts with an inorganic acid (e.g. hydrochloric acid, nitric acid, phosphoric acid, hydrobromic acid, sulfuric acid) or with an organic acid (e.g. acetic acid, formic acid, propionic acid, fumaric acid, maleic acid, succinic acid, tartaric acid, citric acid, maleic acid, ascorbic acid, oxalic acid, benzoic acid, methansulfonic acid, benzenesulfonic acid).

When the object compound (I) has an acid group such as -COOH, the object compound (I) may form a salt with an inorganic base such as sodium, potassium, calcium, magnesium, ammonia, etc. or an organic base such as trimethylamine, etc.


In the following, the method of producing the object compound (I) is described.

The following explanation is applicable not only to the basic compound (I) per se [including a compound usable as the starting compound for preparing another compound included in the definition of compound (I)], but also the salts thereof mentioned above, and, in the following explanation, these compounds are simply referred to as compound (I).

The compound (I) can be produced by allowing, for example, a compound represented by the formula (II):



wherein Z stands for a hydroxyl group or a reactive group of carboxylic acid, and other symbols are of the same meaning as defined above, to react with, for example, a compound represented by the formula (III):



wherein R⁴ is the same as R², except in the case of optionally substituted acyl, namely, a hydrogen atom or an optionally substituted hydrocarbon residue; R³ and n are of the same meaning as defined above, or a salt thereof.

As the reactive group of a carboxylic acid represented by Z, mention is made of halogen (e.g. chlorine, bromine, iodine), a lower(C₁₋₄) alkoxy (e.g. methoxy, ethoxy, propoxy, butoxy) and N-hydroxydiacylimidoester (e.g. N-hydroxysuccinic imidoester, N-hydroxyphthalimidoester, N-hydroxy-5-norbornen-2,3-dicarboxylimidoester), etc.

These reactions are carried out usually in an organic solvent such as hydrocarbon (e.g. pentane, hexane, benzene, toluene), halogenated hydrocarbon (e.g. dichloromethane, chloroform, dichloroethane, carbon tetrachloride), ether (e.g. ethylether, tetrahydrofuran, dioxane, dimethoxyethane), ester (e.g. ethyl acetate, butyl acetate, methyl propionate), amide (e.g. dimethylformamide, dimethylacetamide, hexamethylphosphotriamide), dimethylsulfoxide, etc., under cooling (-10°C to 10°C), at room temperatures (11°C to

40°C), or under heating (41°C to 120°C), and the reaction time ranges usually from 10 minutes to 12 hours. The amount of the compound (III) is preferably 1.0 to 3.0 equivalents relative to the compound (II). This reaction can be carried out, when desired, for example in the case where Z is hydroxy, in the presence of an acid activating agent such as carbonyldiimidazole, dicyclohexylcarbodiimide, diethyl cyanophosphonate, diphenylphosphorylazide, etc., and, in the case where Z is halogen or lower alkoxy, in the presence of an organic base such as pyridine, 4-dimethylaminopyridine, triethylamine, diisopropylamine, triethylenediamine, tetramethylethylenediamine, etc. or in the presence of an inorganic base such as sodium hydrogen carbonate, potassium hydrogen carbonate, lithium hydrogen carbonate, potassium carbonate, sodium carbonate, lithium carbonate, lithium hydroxide, potassium hydroxide, sodium hydroxide, sodium hydride, etc.

In the case where Z is N-hydroxydiacylimidoester, the reaction is carried out in a solvent, for example, dichloromethane, tetrahydrofuran, dioxane, chloroform, dimethylformamide, acetonitrile, water, etc. This reaction is carried out, when necessary, in the presence of an organic or inorganic base. The reaction temperature ranges usually from -10°C to 110°C, preferably from 0°C to 30°C, and the reaction time ranges usually from 5 minutes to 12 hours, preferably from 30 minutes to two hours.

A compound of the formula (II) wherein Z is hydroxy [hereinafter abbreviated as compound (II: Z=hydroxy)], i.e. a free carboxylic acid, can be easily prepared by a per se conventional means, for example, by subjecting the compound (II: Z=lower alkoxy), i.e. the ester compound, to hydrolysis with an alkali metal hydroxide (e.g. potassium hydroxide, lithium hydroxide, sodium hydroxide), an alkali metal carbonate (e.g. potassium carbonate, sodium carbonate, lithium carbonate), mineral acid (e.g. hydrochloric acid, sulfuric acid, nitric acid, phosphoric acid, perchloric acid, hydroiodic acid), an organic acid (e.g. acetic acid, propionic acid, trifluoroacetic acid, monochloroacetic acid, trichloroacetic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid). For the hydrolysis, any conventional solvent can be employed, for example, water, lower(C₁-4 alkanols (e.g. methanol, ethanol, propanol, butanol), dioxane, tetrahydrofuran, dimethylformamide, etc. are preferable. The reaction temperature ranges usually from about -10°C to about 120°C, preferably from 0°C to 80°C. The reaction time ranges usually from 10 minutes to 24 hours, preferably from 30 minutes to 6 hours.

The compound (II: Z=halogen) can be prepared by a per se conventional means, for example, by subjecting carboxylic acid to halogenation using a halogenating agent (e.g. phosphorus oxychloride, phosphorus oxybromide, phosphorus pentachloride, phosphorus pentabromide, thionyl chloride, thionyl bromide, sulfuryl chloride, oxalyl chloride, cyanuric acid chloride, boron tribromide, hydrogen iodide). As acid halides to be obtained by the halogenation are mentioned, for example, acid chloride, acid bromide, acid fluoride and acid iodide, especially acid chloride and acid bromide are preferable.


The above-mentioned halogenation is carried out in the absence of solvent or in a conventional solvent. Preferable solvents are, for example, inactive ones such as chloroform, dichloromethane, dichloroethane, benzene, toluene, etc.

The compound (II: Z=hydroxydiacylimidoester) can be prepared by, in a per se conventional manner, allowing the compound (II: Z=hydroxy) to react with an N-hydroxydicarboxylic acid imide (e.g. N-hydroxysuccinic acid imide, N-hydroxyphthalic acid imide, N-hydroxy-5-norbornen-2,3-dicarboxyimide) in the presence of dicyclohexylcarbodiimide. This reaction is carried out in a conventional solvent (e.g. tetrahydrofuran, dioxane, dimethylformamide, acetonitrile, water, and the compound (II: Z=N-hydroxydiacylimidoester) can be fed to the subsequent reaction without isolation thereof.

The compound (II) usable as the starting material can be prepared by a known method or a method analogous thereto. The compound (III) can also be prepared in accordance with a known method or a method analogous thereto.

Among the compounds (III), for example, the compound (III: R³=CH₂Ph, R⁴=H, n=2) (wherein Ph stands for phenyl group, hereinafter the same abbreviation will be applied.) is a known compound, which is disclosed in Synthesis, 388 (1983).

The compound (III) can be prepared by a per se known method by using a known compound represented by the formula(III'):



wherein R⁶=CO₂C₂H₅, R⁷=CO₂CH₂Ph, and n = an integer of 2 to 6. For example, the compound (III': R⁶=CO₂C₂H₅, R⁷=CO₂CH₂Ph, n = integer of 2 to 6) is subjected to catalytic reduction by a conventional

method or is treated with an acid to give a compound (III': R^5 and n are of the same meaning as defined above, $R^7 = H$), to which was then introduced a hydrocarbon residue by a conventional method to give a compound (III: R^5 and n are of the same meaning as above, $R^7 = R^3$ (R^3 is of the same meaning as above)), then the ester group of R^6 is subjected to amidation directly by a conventional means or amidation after
 5 converting the ester group of R^6 into a carboxyl group to thereby obtain a compound (III': $R^6 = CONHR^4$, $R^7 = R^3$ (R^3 and n are of the same meaning as defined above)), followed by subjecting the compound to reduction with lithium aluminium hydride by per se conventional means to afford the compound (III).

The above-mentioned known compound, the starting compound, (III': $R^6 = CO_2CH_2H_5$, $R^7 = CO_2CH_2Ph$, $n =$ an integer of 2 to 6) is that disclosed in Japanese Patent Publication (laid open) 99476/1981 and
 10 Chem. Pharm. Bull. 34, 3747(1986).

Among the object compounds (I) of the present invention, a compound (I) in which $R^5 = R^2$ (except for the case of $R^5 =$ hydrogen, wherein R^5 is an optionally substituted hydrocarbon residue or acyl group) can also be prepared by introducing hydrocarbon residue into the object compound of this invention (I: $R^2 = H$) or by subjecting the said compound to acylation. For example, the above compound (I: $R^2 = R^5$) can be
 15 prepared also by allowing a compound (I: $R^2 = H$) to react with a compound represented by the formula (IV): R^5-Y (wherein R^5 is defined as above; when R^5 is an optionally substituted hydrocarbon residue, Y stands for halogen, and, when R^5 is an optionally substituted acyl group, Y stands for hydroxy, OR^5 or a reactive group of carboxylic acid) by a per se known method.

In the reaction of the compound (I: $R^2 = H$) with a compound (IV), use of a solvent is not always
 20 required, and, when required, use of an organic solvent such as hydrocarbon (e.g. pentane, hexane, benzene, toluene), halogenated hydrocarbon (e.g. dichloromethane, chloroform, dichloroethane, carbon tetrachloride), ether (e.g. ethyl ether, tetrahydrofuran, dioxane, dimethoxyethane), amide (e.g. dimethylformamide, hexamethylphosphotriamide), dimethylsulfoxide, etc. is preferable. The reaction is carried out at temperatures ranging from $-10^\circ C$ to $200^\circ C$, preferably from $0^\circ C$ to $120^\circ C$. The reaction time ranges
 25 usually from 5 minutes to 12 hours, preferably from 10 minutes to 6 hours. The amount of the above-mentioned compound (IV) is usually equimolar or in an excess relative to the compound (I: $R^2 = H$), preferably 1.1 to 20.0 times as much per mole. When R^5 is an optionally substituted hydrocarbon residue and Y is halogen, the reaction is carried out in the presence of, for example, an organic base such as pyridine, 4-dimethylaminopyridine, triethylamine, diisopropylamine, triethylenediamine,
 30 tetramethylethylenediamine, etc. or an inorganic base such as sodium hydride, metallic sodium, potassium amide, sodium hydrogen carbonate, potassium hydrogen carbonate, sodium carbonate, potassium carbonate, lithium hydroxide, potassium hydroxide, sodium hydroxide, etc. The amount of these bases is generally equimolar to excess relative to the compound (I: $R^2 = H$), preferably 1.1 to 5 times as much mol.

And, Examples of the reactive groups of carboxylic acid represented by Y when R^5 stands for an acyl
 35 group include halogen (e.g. chlorine, bromine, iodine), lower (C_1-4) alkoxy (e.g. methoxy, ethoxy, propoxy, butoxy) and N-hydroxydiacylimidoester (e.g. N-hydroxysuccinic acid imidoester, N-hydroxyphthalic acid imidoester, N-hydroxy-5-norbornene-2,3-dicarboxylimidoester), etc.

And, the reaction of the compound (I: $R^2 = H$) with a compound (IV), when R^5 is an acyl group, can be carried out, if desired, when Y is hydroxy, in the presence of an acid-activating agent such as carbonyl-
 40 diimidazole, dicyclohexylcarbodiimide, diethyl cyanophosphate, diphenylphosphorylazide, etc., when Y is OR^5 , in the presence of a mineral acid such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, etc., an organic acid such as acetic acid, formic acid, propionic acid, methanesulfonic acid, p-toluenesulfonic acid, etc. or an acyl halogenide whose acyl group is the same as R^5 , and, when Y is halogen or a lower alkoxy, in the presence of an organic base such as pyridine, 4-dimethylaminopyridine,
 45 triethylamine, diisopropylamine, triethylenediamine, tetramethylethylenediamine, etc. or an inorganic base such as sodium hydrogen carbonate, potassium hydrogen carbonate, lithium hydrogen carbonate, lithium hydroxide, potassium hydroxide, sodium hydroxide, etc.

Further, when Y is a N-hydroxydiacylimido ester, the reaction is carried out in a solvent, for example, preferably dichloromethane, tetrahydrofuran, dioxane, chloroform, dimethylformamide, acetonitrile, water,
 50 etc. This reaction can be carried out, upon desire, in the presence of an organic or an inorganic base mentioned above referring to the case where Y is halogen or lower alkoxy.

When the reaction is carried out in the presence of the above-mentioned acid-activating agent, acid, halide and base, the amount of these agents ranges generally from equimolar to excess relative to the compound (I: $R^2 = H$), preferably in 1.1 to 5 times molar excess.

And, among the object compounds (I) of the present invention, the compound (I: $X = NH_2$) can also be prepared by subjecting also the object compound of the present invention (I: $X = NO_2$) or a salt thereof. The reduction can be conducted by a per se known method, for example those disclosed in J. Am. Chem. Soc.,
 55 49 1093(1927) or Ber., 76, 1011(1943) or an equivalent.

This reaction can be carried out, for example, by conducting a catalytic reduction in hydrogen streams, in the presence of a catalyst (e.g. palladium-carbon, platinum dioxide) at normal temperatures under normal pressure. As the solvent, use is made of, for example, methanol, ethanol, water, dimethylformamide, dioxane, etc., but any other solvents can be used so long as they do not inhibit this reaction. This reaction can be carried out, when desired, in the presence of a mineral acid such as hydrochloric acid, hydrobromic acid, sulfuric acid, etc. or an organic acid such as acetic acid, formic acid, propionic acid, oxalic acid, etc.

A compound (I), wherein X stands for an acylamino group (e.g. acetylamino, benzoylamino, benzenesulfonylamino) can be prepared by subjecting a compound (I: $X = NH_2$) to acylation. This acylation can be carried out by allowing the compound (I: $X = NH_2$) to react with an acylating agent, for example, an acid (e.g. acetic acid, propionic acid, benzoic acid, benzenesulfonic acid, p-toluenesulfonic acid), C_1 - ∞ alkyl ester of an acid (e.g. methyl acetate, ethyl propionate, methyl benzenesulfonate), an acid halogenide (e.g. acetyl chloride, acetyl bromide, p-toluenesulfonic acid chloride, benzenesulfonyl chloride), an acid anhydride (acetic anhydride, propionic anhydride, benzoic anhydride) or N-hydroxydiacylimido ester of an acid (e.g. N-acetyloxysuccinimide, N-benzoyloxypthalimide, N-acetyloxy-5-norbornene-2,3-dicarboxyimide), etc.

These reactions can usually be carried out in an organic solvent such as hydrocarbon (e.g. pentane, hexane, benzene, toluene), halogenated hydrocarbon (e.g. dichloromethane, chloroform, dichloroethane, carbon tetrachloride), ether (e.g. ethyl ether, tetrahydrofuran, dioxane, dimethoxyethane), ester (e.g. ethyl acetate, butyl acetate, methyl propionate), amide (e.g. dimethylformamide, dimethylacetamide, hexamethylphosphotriamide), dimethylsulfoxide, etc., under cooling ($-10^\circ C$ to $10^\circ C$), at room temperatures ($11^\circ C$ to $40^\circ C$) or under heating ($41^\circ C$ to $120^\circ C$), and the reaction time ranges from 10 minutes to 12 hours. The amount of the above-mentioned acylating agents ranges preferably from 1.0 to 3.0 equivalents relative to the compound (I: $X = NH_2$). Further, this reaction can be carried out, if desired, when the acylating agent is an acid, in the presence of, for example, an acid-activating agent such as carbonyldiimidazole, dicyclohexylcarbodiimide, diethyl cyanophosphate, diphenylphosphorylazide, etc., and, if the acylating agent is C_1 - ∞ alkyl ester of an acid or an acid halide, in the presence of an organic base such as pyridine, 4-dimethylaminopyridine, triethylamine, diisopropylamine, triethylenamine, tetramethylethylenediamine, etc. or in the presence of an inorganic base such as sodium hydrogen carbonate, potassium hydrogen carbonate, lithium carbonate, lithium hydroxide, potassium hydroxide, sodium hydroxide, etc., and, if the acylating agent is an acid anhydride, in the presence of a mineral acid such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, etc. or an organic acid such as acetic acid, formic acid, propionic acid, methanesulfonic acid, p-toluenesulfonic acid, etc.

Further, if the acylating agent is an N-hydroxydiacylimido ester, the acylation is carried out preferably in the presence of a solvent such as dichloromethane, tetrahydrofuran, dioxane, chloroform, dimethylformamide, acetonitrile, water, etc. This reaction can be carried out, when desired, in the presence of such an organic or inorganic base as mentioned above. The reaction temperature ranges, usually, from $-10^\circ C$ to $110^\circ C$, preferably from $0^\circ C$ to $30^\circ C$, and the reaction time ranges, usually, from 5 minutes to 12 hours, preferably from 30 minutes to 2 hours.

The compound (I) of the present invention acts on the central nervous system of mammals, has a strong anticholinesterase activity and shows an excellent antiamnesic action against various types of induction of amnesia in humans and animals such as mice.

The compound (I) of the present invention has, as compared with physostigmine, such characteristic features as showing a very good separability of the action on the central nerves from that on peripheral nerves, being accompanied with no or very slight actions on peripheral nerves causing convulsion, salivation, diarrhea, in the dosage showing antiamnesic action. The compound (I) has prolonged action and is of low toxicity; it is a remarkably effective by oral administration.

Therefore, the present compounds are useful in agents for improving cerebral functions in a mammal including human beings.

The diseases to which the compounds of the present invention can be effectively applied are senile dementia, Alzheimer's disease, Huntington's chorea, hyperkinesia, mania, etc., and the compounds can be used for the prophylaxis or therapy of these diseases.

The compounds of the present invention can be administered orally or non-orally to mammals including man in various dosage forms such as tablets, granules, capsules, injections, suppositories and so forth. The dose varies with the kinds of diseases, symptoms, etc. Generally, however, in the case of oral administration, the daily dose ranges from 0.001 mg to 100 mg, preferably from 0.01 mg to 10 mg.

The following examples, reference examples, formulation examples and experimental examples are intended to illustrate the present invention in further detail and should by no means be construed as limiting the scope of the present invention.


The elution in column chromatography in the examples and reference examples were conducted,

unless otherwise specified, using the technique of Thin Layer Chromatography (TLC). The eluate fractions containing the object compound were confirmed and collected by employing, as a supplemental means of detection, the procedure which comprises spraying 48% HBr onto the spot on the TLC plate, hydrolyzing by heating, then spraying thereon a ninhydrin reagent, and heating again to cause change of the color to red to reddish purple. Unless otherwise specified, the silica gel used for the column was Kiesel-gel 60 (70 to 230 mesh) manufactured by E. Merck AG.

The term "room temperatures" usually ranges from about 5° C to 40° C.
Unless otherwise specified, % means weight percentage.

Example 1

(E)-3-Phenyl-N-[2-(1-benzylpiperidin-4-yl)ethyl]-2-propenamide*hydrochloride




To a dimethylformamide solution (20 ml) containing (E)-cinnamic acid (1.05 g), 4-(2-aminoethyl)-1-benzylpiperidine*dihydrochloride (1.8 g) and triethylamine (1.0 ml) was added, under ice-cooling, diethyl cyanophosphonate (1.7 g). The mixture was stirred for one hour under ice-cooling, and there was added water (100 ml). The mixture was subjected to extraction with dichloromethane. The dichloromethane solution was dried over anhydrous sodium sulfate, then the solvent was distilled off under reduced pressure. The oily residue was subjected to a silica gel column chromatography [developing solvent : ethyl acetate - methanol = 20 : 1(V/V)]. The solvent of the solution containing the object product was distilled off. To the residue was added an ethanolic solution (2.4 ml) of 3N hydrochloric acid, then the solvent was distilled off. The residual solid was recrystallized from ethanol-ether [5:1(V/V)] to give colorless crystals (1.2 g) m.p. 125 to 127° C.

Elemental Analysis for $C_{23}H_{28}N_2O \cdot HCl$:

Calcd. :	C 71.76	H 7.59	N 7.28
Found :	C 71.62	H 7.49	N 7.01

Example 2

(E)-3-Phenyl-N-methyl-N-[2-(1-benzylpiperidin-4-yl)ethyl]-2-propenamide*hydrochloride



To a dimethylformamide solution (5 ml) of (E)-3-phenyl-N-[2-(1-benzylpiperidin-4-yl)ethyl]-2-propenamide*hydrochloride (0.6 g) obtained in Example 1 was gradually added sodium hydride (80 mg) at room temperature and the mixture was stirred at 60° C for 30 minutes. The reaction mixture was cooled on an ice-bath and there was added methyl iodide (0.21 g), followed by stirring at room temperature for one


hour. To the resultant mixture was added water and the mixture was subjected to extraction with dichloromethane. The dichloromethane solution was washed with water and dried over anhydrous sodium sulfate, then the solvent was distilled off. The oily residue was subjected to a silica gel column chromatography [developing solvent : ethyl acetate - methanol = 10 : 1(V/V)]. Then the solvent of the solution containing the object compound was distilled off. To the residue was added an ethanolic solution (0.5 ml) of 3N hydrogen chloride, followed by distilling off the solvent to obtain an amorphous powder (0.2 g).

Elemental Analysis for $C_{24}H_{30}N_2O \cdot HCl$:

Calcd. :	C 72.25	H 7.83	N 7.02
Found :	C 72.01	H 7.79	N 6.95

Example 3

(E)-3-Phenyl-N-acetyl-N-[2-(1-benzylpiperidin-4-yl)ethyl]-2-propenamide*hydrochloride



(A) A mixture of (E)-3-phenyl-N-[2-(1-benzylpiperidin-4-yl)ethyl]-2-propenamide*hydrochloride (0.5 g), acetic anhydride (2.5 ml) and a catalytic amount of p-toluenesulfonic acid*monohydrate was stirred at 80 °C for 6 hours. The reaction mixture was left standing for cooling, after there was added water. To the mixture was added 10% NaOH to render the pH to about 9 to 10, followed by extraction with dichloromethane. The dichloromethane solution was washed with water and dried over anhydrous sodium sulfate, followed by distilling off the solvent. The oily residue was subjected to a silica gel column chromatography (developing solvent : ethyl acetate). The solvent of the solution containing the object compound was distilled off. To the residue was added an ethanol solution (0.44 ml) of 3N hydrogen chloride, and the solvent was distilled off to obtain a hygroscopic amorphous powder (0.45 g).

Elemental Analysis for $C_{25}H_{30}N_2O_2 \cdot HCl$:

Calcd. :	C 70.32	H 7.32	N 6.56
Found :	C 70.18	H 7.29	N 6.41

(B) A pyridine solution (5 ml) of (E)-3-phenyl-N-[2-(1-benzylpiperidin-4-yl)ethyl]-2-propenamide*hydrochloride (0.25 g) and acetyl chloride (0.3 ml) was stirred at 60 °C for 2 hours. Water was added to the reaction mixture after being left standing for cooling; the mixture was dried over anhydrous sodium sulfate. The solvent was distilled off. The oily residue was subjected to silica gel column chromatography (developing solvent : ethyl acetate). The solvent of the solution containing the object compound was distilled off. To the residue was added an ethanol solution of 3N hydrogen chloride. The solvent was then distilled off to obtain hygroscopic amorphous powder (0.16 g).


Elemental Analysis for $C_{25}H_{30}N_2O_2 \cdot HCl$:







Calcd. :	C 70.32	H 7.32	N 6.56
Found :	C 70.26	H 7.13	N 6.48









Example 4

By the procedure of Example 1, compounds as set forth in Table 1 were obtained.

Table 1




Compound No.		R ¹	n	m.p. (°C)	Molecular Formula	Elemental Analysis Calcd. (Found)		
						C	H	N
1		CH ₃	2	amorphous	C ₂₄ H ₃₀ N ₂ O·HCl	72.25 (72.18)	7.83 7.74	7.02 6.98
2		C ₆ H ₅	2	amorphous	C ₂₉ H ₃₂ N ₂ O·HCl	75.55 (75.48)	7.21 7.11	6.08 6.00
3		H	2	113 - 115	C ₂₉ H ₂₇ N ₃ O	75.61 (75.52)	7.79 7.72	12.02 11.96
4		H	2	180 - 183	C ₂₁ H ₂₆ N ₂ O ₂ ·HCl	67.28 (67.27)	7.26 7.11	7.47 7.39
5		H	2	89 - 92	C ₂₁ H ₂₆ N ₂ OS·HCl	64.51 (64.44)	6.96 6.93	7.17 7.01

Compound No.		R ¹	n	m.p. (°C)	Molecular Formula	Elemental Analysis		
						Calcd.	(Found)	
						C	H	N
6		H	2	amorphous	C ₂₅ H ₃₂ N ₂ O ₃ ·HCl	67.48 (67.32)	7.47 7.45	6.30 6.27)
7			2	amorphous	C ₂₇ H ₃₄ N ₂ O ₃ ·HCl	68.85 (68.81)	7.49 7.45	5.95 5.86)
8		H	2	amorphous	C ₂₃ H ₂₇ N ₃ O ₃ ·HCl	64.25 (64.11)	6.56 6.42	9.77 9.61)
9		H	2	amorphous	C ₂₄ H ₂₇ N ₃ O·HCl	70.32 (70.14)	6.88 6.83	10.25 10.01)
10		H	2	158 - 160	C ₂₄ H ₃₀ N ₂ O ₃ S	67.58 (67.39)	7.09 6.98	6.57 6.43)
11		H	2	amorphous	C ₂₄ H ₃₀ N ₂ O·HCl	72.25 (72.06)	7.83 7.77	7.02 6.93)
12		H	2	108 - 110	C ₂₃ H ₂₇ ClN ₂ O	72.14 (72.03)	7.11 6.99	7.32 7.03)

Example 5

By a procedure similar to that of Example 3, compounds as set forth in Table 2 were obtained.

Table 2



Compound No.		R1	n	m.p. (°C)	Molecular Formula	Elemental Analysis Calcd. (Found)		
						C	H	N
1		CH ₃	2	amorphous	C ₂₆ H ₃₂ N ₂ O ₂ ·HCl	70.81 (70.67)	7.54 7.50	6.35 6.29)
2		C ₆ H ₅	2	amorphous	C ₃₇ H ₃₄ N ₂ O ₂ ·HCl	74.01 (73.94)	7.01 6.99	5.57 5.54)

Formulation Example 1

(1) (E)-3-phenyl-N-[2-(1-benzylpiperidin-4-yl)ethyl]-2-propenamide•hydrochloride	1 g
(2) lactose	197 g
(3) corn starch	50 g
(4) magnesium stearate	2 g

The above ingredients (1), (2) and corn starch (20 g) were mixed. The mixture was granulated with a paste from corn starch (15 g) and water (25 ml). To the granules was added corn starch (15 g), and the mixture was compressed with a tableting machine into 2000 tablets (3 mm diameter), each containing 0.5 mg of (1).

Formulation Example 2

(1) (E)-3-phenyl-N-[2-(1-benzylpiperidin-4-yl)ethyl]-2-propenamide•hydrochloride	2 g
(2) lactose	196 g
(3) corn starch	50 g
(4) magnesium stearate	2 g

The above ingredients (1), (2) and corn starch (20 g) were mixed. The mixture was granulated with a paste prepared from corn starch (20 g) and water (25 ml). To the granules were added corn starch (15 g) and (4), and the mixture was compressed into 2000 tablets (5 mm diameter), each containing 1 mg of (1).

Experimental Example

Cholinesterase activity was measured radiometrically by the method of Johnson and Russell ¹⁾, modified by Kleinberger and Yanai ²⁾, with a slight modification.

S₁ fraction of the cerebral cortex of male Wistar rats, the enzyme source, was preincubated in a scintillation vial with drugs for 15 min at room temperature, and then [acetyl-³H]-acetylcholine (final 200 μ M) was added and incubated further for 30 min. The reaction was terminated by adding solution containing 1 M chloroacetic acid, followed by toluene-based scintillant, and the vials were capped and shaken to transfer the product, [³H]-acetic acid, to toluene phase. Then radioactivity in the toluene phase was counted by liquid scintillation spectrometry. Inhibitory activity of the test drug was expressed by 50%-inhibitory concentration (IC₅₀), which was calculated by linear regression of log-probit transformation of the inhibition curve. By the same method, cholinesterase activity of physostigmine was measured.

The results were shown.

1) C. D. Johnson and R. L. Russell (1975) Anal. Biochem. 64, 229-238.


2) N. Kleinberger and J. Yanai (1985) Dev. Brain Res. 22, 113-123.

Table 3

Compound (Example No.)	Anti-acetylcholinesterase activity IC ₅₀ (μM)
1	9.6
2	18
3	0.64
4 - 1	8.1
4 - 2	3.5
4 - 3	2.0
4 - 4	9.9
4 - 5	13
4 - 6	2.6
4 - 7	1.2
4 - 8	2.8
4 - 9	3.3
4 - 10	1.6
4 - 11	15
4 - 12	9.2
physostigmine	0.22

Claims

1. A compound of the formula:



wherein ring A stands for an optionally substituted aromatic ring; R¹ stands for a hydrogen atom or an optionally substituted hydrocarbon residue or forms an optionally substituted carbocyclic ring with the adjacent group -CH=C- together with two carbon atoms constituting the ring A; R² stands for a hydrogen atom, an optionally substituted hydrocarbon residue or an optionally substituted acyl group; R³ stands for an optionally substituted hydrocarbon residue; and n denotes an integer ranging from 2 to 6, or salts thereof.

2. A compound according to claim 1, wherein ring A is aromatic monocyclic hydrocarbon, aromatic condensed polycyclic hydrocarbon or aromatic heterocyclic ring.

3. A compound according to claim 2, wherein the aromatic monocyclic hydrocarbon is benzene.

4. A compound according to claim 2, wherein the aromatic condensed polycyclic hydrocarbon is naphthalene or anthracene.

5. A compound according to claim 2, wherein the aromatic heterocyclic ring is a member selected from the group consisting of thiophene, furan, pyrazole, thiazole, isothiazole, oxazole, isoxazole, imidazole, triazole, tetrazole, pyridine, pyrimidine and pyridazine.

6. A compound according to claim 2, wherein the ring A is unsubstituted or substituted with 1 to 3 of C₁₋₄ alkyl, halogen, nitro, cyano, acetylamino, C₁₋₄ alkoxy, optionally substituted phenyl, optionally substituted benzyl, optionally substituted benzoyl, optionally substituted benzoylamino, optionally substituted C₁₋₆ alkylsulfonyl, optionally substituted benzylsulfonyl, optionally substituted phenylsulfonylamino, optionally substituted benzylsulfonylamino, optionally substituted phenylcarbamoyl, methoxycarbonyl or/and diethoxycarbonyl.

7. A compound according to claim 1, wherein R¹ is hydrogen, C₁₋₁₁ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C₃₋₇ monocyclic cycloalkyl, C₈₋₁₄ cross-linked cyclic saturated hydrocarbon or cyclic unsaturated hydrocarbon residue.

8. A compound according to claim 1, wherein R¹ form a 5- to 7-membered ring together with the adjacent group -CH=C- and two carbon atoms constituting the ring A.

9. A compound according to claim 8, wherein the condensed ring formed by the 5- to 7-membered ring and ring A is unsubstituted or substituted with 1 to 3 of C₁₋₄ alkyl, halogen, C₁₋₄ alkoxy, amino, mono- or di-C₁₋₄ alkyl-substituted amino, nitro, cyano or/and C₁₋₄ alkoxycarbonyl.

10. A compound according to claim 1, wherein R² is hydrogen, C₁₋₁₁ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C₃₋₇ monocyclic cycloalkyl, C₈₋₁₄ cross-linked cyclic saturated hydrocarbon, cyclic unsaturated hydrocarbon residue, carboxylic acid acyl, carbamic acid acyl, sulfonic acid acyl, substituted hydroxycarboxylic acid acyl.

11. A compound according to claim 1, wherein R³ is C₁₋₃ alkyl-C₆₋₁₀ aromatic hydrocarbon.

12. A compound according to claim 1, wherein n is an integer of 2, 3 or 4.

13. A compound according to claim 1, wherein ring A is benzene, pyridine, furan or thiophene which may be substituted with C₁₋₄ alkyl, C₁₋₄ alkoxy, nitro, cyano, halogen or/and C₁₋₆ alkylsulfonyl; R¹ is hydrogen, C₁₋₆ alkyl or phenyl, or forms 1,2-dihydronaphthalene which may be substituted with C₁₋₄ alkoxy together with the adjacent group -CH=C- and two carbon atoms of the ring A; R² is hydrogen, C₁₋₆ alkyl or C₁₋₆ alkylcarbonyl; R³ is benzyl; and n is 2.

14. A compound according to claim 1, wherein ring A is benzene, pyridine, furan or thiophene which may be substituted with C₁₋₄ alkoxy; R¹ is hydrogen, C₁₋₆ alkyl or phenyl, or forms 1,2-dihydronaphthalene which may be substituted with C₁₋₄ alkoxy together with the adjacent group -CH=C- and two carbon atoms of ring a; R² is hydrogen, C₁₋₆ alkyl or C₁₋₆ alkylcarbonyl; R³ is benzyl, and n is 2.


15. A compound according to claim 1, wherein ring A is benzene; R¹ is hydrogen; R² is hydrogen, C₁₋₆ alkyl or C₁₋₆ alkylcarbonyl; R³ is benzyl; and n is 2.

16. A compound according to claim 1, which is (E)-3-phenyl-N-acetyl-N-[2-(1-benzylpiperidin-4-yl)ethyl]-2-propenamide*hydrochloride.

17. A compound according to claim 1, which is (E)-3-(3-pyridyl)-N-[2-(1-benzylpiperidin-4-yl)ethyl]-2-propenamide.


18. A compound according to claim 1, which is 3,4-dihydro-6,7-dimethoxy-N-[2-(1-benzylpiperidin-4-yl)ethyl]naphthalene-2-carboxamide*hydrochloride

19. A method of preparing a compound according to claim 1, which is characterized by (A) allowing a compound represented by a compound represented by the formula:




wherein Z stands for a hydroxyl group or a reactive group of carboxylic acid, and other symbols are of the same meaning as defined in Claim 1,

to react with a compound represented by the formula:




wherein R⁴ stands for a hydrogen atom or an optionally substituted hydrocarbon residue; R³ and n are of the same meaning as defined in Claim 1 or a salt thereof to give an unsaturated carboxylic acid amide derivative represented by the formula:




wherein each symbol is of the same meaning as defined above, or a salt thereof,

(B) subjecting a compound represented by the formula:




wherein each symbol is of the same meaning as defined above, to a reaction introducing a hydrocarbon residue or acylation to give an unsaturated carboxylic acid amide derivative represented by the formula:




wherein R^5 stands for an optionally substituted hydrocarbon residue or acyl group, and the remaining symbols are of the same meaning as defined above, or a salt thereof,

(C) subjecting a compound represented by the formula:




wherein R^2 is as defined in Claim 1, and the remaining symbols are of the same meaning as above, or a salt thereof to reduction to give an unsaturated carboxylic acid amide derivative of the formula:




wherein each symbol is of the same meaning as above, or a salt thereof, or

(D) subjecting a compound represented by the formula:



wherein each symbol is of the same meaning as defined above, or a salt thereof to acylation to give an

unsaturated carboxylic acid amide derivative represented by the formula:



10 wherein X' stands for acylamino group, and the remaining symbols are of the same meaning as defined above, or a salt thereof.

20. A pharmaceutical composition which contains a compound according to claim 1 together with carrier, diluent therefor.

21. A method for improving cerebral functions in a mammal which comprises administering to the
15 mammal an effective amount of a compound according to claim 1.

20

25

30

35

40

45

50

55



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT
which under Rule 45 of the European Patent Convention
shall be considered, for the purposes of subsequent
proceedings, as the European search report

Application number

EP 89 10 2376

DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
A	EP-A-0 229 391 (EISAI) * Claims 1, 25 * -----	1, 21	C 07 D 211/26 C 07 D 401/12 C 07 D 405/12 C 07 D 409/12 A 61 K 31/445

**TECHNICAL FIELDS
SEARCHED (Int. Cl.4)**

C 07 D 211/00
C 07 D 401/00
C 07 D 405/00
C 07 D 409/00
A 61 K 31/00

INCOMPLETE SEARCH

The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.

Claims searched completely: 1-20

Claims searched incompletely:

Claims not searched: 21

Reason for the limitation of the search:

Method for treatment of the human or animal
body by surgery or therapy (see art. 52(4)
of the European Patent Convention).

Place of search
The Hague

Date of completion of the search
17-05-1989

Examiner
CASADO MARTI DE MERCADO

CATEGORY OF CITED DOCUMENTS

X : particularly relevant if taken alone
Y : particularly relevant if combined with another
document of the same category
A : technological background
O : non-written disclosure
P : intermediate document

T : theory or principle underlying the invention
E : earlier patent document, but published on, or
after the filing date
D : document cited in the application
L : document cited for other reasons

& : member of the same patent family, corresponding
document